

2020
MATHEMATICS
[HONOURS]
Paper : II

Full Marks : 100

Time : 4 Hours

*The figures in the right-hand margin indicate marks.**Symbols, notations have their usual meanings.***GROUP-A****(Differential Calculus)****[Marks : 35]**1. Answer any **three** questions: 1×3=3

a) If $u = xyf\left(\frac{y}{x}\right)$, then find $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$.

b) Find $\lim_{x \rightarrow 0^+} \frac{1}{e^x + 1}$.

c) Evaluate the following limit (if exist):

$$\lim_{x \rightarrow 0} \frac{3x + |x|}{7x - 5|x|}.$$

d) Find the nature of discontinuity of the function

$$f(x) = \begin{cases} (x+1)\sin\frac{1}{x} & , x \in (-1, 1) \\ 0 & , \text{ otherwise} \end{cases}$$

at $x=1$.

e) Give an example of a function which is continuously at a point but not differentiable at that point.

2. Answer any **two** questions: 2×2=4

a) If $y = x^{n-1} \log x$ then prove that $y_n = \frac{(n-1)!}{x}$.

b) Prove that the curve $r^n = a^n \cos n\theta$ and $r^n = b^n \sin n\theta$ cut orthogonally.c) Find the derived function f' corresponding to the function $f : [0, 3] \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} x & , \text{ when } 0 \leq x \leq 1 \\ 2 - x^2 & , \text{ when } 1 < x < 2 \\ x - x^2 & , \text{ when } 2 \leq x \leq 3. \end{cases}$$

3. Answer any **three** questions: $6 \times 3 = 18$

a) Let $f : [a, b] \rightarrow \mathbb{R}$ be such that $f''(x)$ exists in $[a, b]$ and $f'(a) = f'(b)$. Prove that

$$f\left(\frac{a+b}{2}\right) = \frac{1}{2}[f(a) + f(b)] + \frac{1}{8}(b-a)^2 f''(c)$$

for some $c \in (a, b)$. 6

b) i) Find all asymptotes of the following curve:

$$y = \frac{x^2 - x - 2}{x - 2}$$

ii) Find a and b such that

$$\lim_{x \rightarrow 0} \frac{ae^x + be^{-x} + 2 \sin x}{\sin x + x \cos x} = 2. \quad 3+3$$

c) If $y = \log\left[x + \sqrt{1+x^2}\right]$, then prove that

$$y_{2n}(0) = 0 \text{ and } y_{2n+1}(0) = (-1)^n \cdot 1^2 \cdot 3^2 \cdot 5^2 \dots (2n-1)^2. \quad 6$$

d) Define pedal equation of a curve. Show that the pedal equation of the curve

$$r \cos\left(\frac{\sqrt{a^2 - b^2}}{a} \theta\right) = \sqrt{a^2 - b^2} \text{ is } p\sqrt{b^2 + r^2} = ar. \quad 1+5$$

e) i) Find $\frac{\partial z}{\partial x}$ for the following function:

$$x^2 \sin(2y - 5z) = 1 + y \cos(6zx)$$

ii) A function f is defined on $[0, 1]$ by $f(0) = 1$ and

$f(x) = 0$ if x be irrational

$$= \frac{1}{n}, \text{ if } x = \frac{m}{n} \text{ where } m, n \text{ are}$$

positive integers prime to each other.

Prove that f is continuous at every irrational point in $[0, 1]$ and discontinuous at every rational point in $[0, 1]$. 2+4

4. Answer any **one** question: $10 \times 1 = 10$

a) i) If ρ_1, ρ_2 be the radii of curvature at the end of a focal chord of the parabola $y^2 = 4ax$ then show that

$$\rho_1^{-\frac{2}{3}} + \rho_2^{-\frac{2}{3}} = (2a)^{-\frac{2}{3}}.$$

ii) Find the multiple points of the curve

$$x^4 - 4ax^3 - 4ay^3 + 4a^2x^2 + 3a^2y^2 - a^4 = 0.$$

iii) Discuss the nature of discontinuity at $x=1$ of the function

$$f(x) = \lim_{n \rightarrow \infty} \frac{\log(2+x) - x^{2n} \sin x}{1+x^{2n}}.$$

- iv) If f is monotonically increasing function on $[a, b]$ and $a < c < b$, then show that $f(c+0)$ exists. 3+2+2+3=10

- b) i) If $x^2 + y^2 + z^2 - 2xyz = 1$ then show that

$$\frac{dx}{\sqrt{1-x^2}} + \frac{dy}{\sqrt{1-y^2}} + \frac{dz}{\sqrt{1-z^2}} = 0.$$

- ii) If $u = \log(x^3 + y^3 + z^3 - 3xyz)$ then show that $u_{xx} + u_{yy} + u_{zz} = -\frac{3}{(x+y+z)^2}$.

- iii) If $f(x) = \begin{cases} 1 & \text{when } x \text{ is rational} \\ 0 & \text{when } x \text{ is irrational,} \end{cases}$

state with reasons, which of the following statement is true:

- p. f is continuous at rational points, but discontinuous at irrational points.
 q. f is continuous at irrational points and discontinuous at rational points.
 r. f is continuous every where
 s. f is discontinuous everywhere.

3+3+4=10

GROUP-B
(Integral Calculus)

[Marks : 25]

5. Evaluate any **one** of the following: 3×1=3

a) $\int \frac{dx}{(1-x)\sqrt{1-x^2}}$

b) $\int \sin^{-1}\left(\sqrt{\frac{x}{a+x}}\right) dx$

6. Answer any **two** questions: 6×2=12

- a) Prove that

$$\int_0^{\frac{\pi}{2}} \frac{dx}{(a^2 \cos^2 x + b^2 \sin^2 x)^2} = \frac{\pi}{4} \cdot \frac{a^2 + b^2}{a^3 \cdot b^3}; \quad a, b > 0.$$

6

- b) i) Find

$$\lim_{n \rightarrow \infty} \left[\frac{n}{n^2} + \frac{n}{1^2 + n^2} + \frac{n}{2^2 + n^2} + \dots + \frac{n}{(n-1)^2 + n^2} \right].$$

- ii) Evaluate: $\int (\sin^{-1} x)^4 dx$. 3+3

c) Evaluate: 3+3

i) $\int \frac{dx}{\sin(x-a)\sin(x-b)}$

ii) $\int \frac{x^2 + 2x + 3}{\sqrt{1-x^2}} dx$

7. Answer any **one** question: 10×1=10

a) i) Find the moment of inertia of a thin uniform lamina in the form of an ellipse

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1 \text{ about its axes.}$$

ii) Find the volume of the solid obtained by the revolution of the cissoid $y^2(2a-x) = x^3$ about its asymptote.

iii) If $u_n = \int_0^1 x^n \tan^{-1} x \, dx$, prove that for $n > 2$,

$$(n+1)u_n + (n-1)u_{n-2} = \frac{\pi}{2} - \frac{1}{n}. \quad 3+4+3$$

b) i) If $I_n = \int_0^{\frac{\pi}{2}} x^n \sin x \, dx$, n being a positive integer greater than 1, then show that

$$I_n + n(n-1)I_{n-2} = n\left(\frac{\pi}{2}\right)^{n-1}.$$

Hence find the value of $\int_0^{\frac{\pi}{2}} x^5 \sin x \, dx$.

ii) Find the volume of the solid generated by revolving the cardioide $r = a(1 - \cos\theta)$ about the initial line. 6+4

GROUP-C

(Differential Equation-I)

[Marks : 40]

8. Answer any **two** questions: 1×2=2

a) Determine the order and degree of the differential equation

$$\left\{ 1 + \left(\frac{dy}{dx} \right)^2 \right\}^{\frac{3}{2}} = \frac{1}{1 + \frac{dy}{dx}}.$$

b) Check whether the differential equation

$$(x^2 - 2xy - y^2)dx - (x + y)^2 dy = 0$$

is exact.

c) Determine Wronskian of the functions

$$e^x, e^{2x}, e^{3x}.$$

9. Answer any **two** questions: $2 \times 2 = 4$

a) Find the orthogonal trajectories of the family of curves $xy = a^2$.

b) Solve the equation: $x dy - y dx - \cos \frac{1}{x} dx = 0$.

c) Show that the substitution $x = \sinh z$ transforms the equation $(1+x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} = 4y$ into $\frac{d^2y}{dz^2} = 4y$.

10. Answer any **four** questions: $6 \times 4 = 24$

a) Solve the following differential equation

$$(x+a)^2 \frac{d^2y}{dx^2} - 4(x+a) \frac{dy}{dx} + 6y = x.$$

b) Find the eigen values and eigen functions of the equation

$$\frac{d^2y}{dx^2} + \lambda y = 0, \quad 0 \leq x \leq \pi$$

satisfying the boundary conditions $y = 0$ at

$$x = 0 \text{ and } \frac{dy}{dx} = 0 \text{ at } x = \pi.$$

c) Show that the system of co-axial parabolas $y^2 = 4a(x+a)$ is self orthogonal.

d) Reduce the differential equation $y = 2px - p^2y$ to Clairant's form by the substitutions $y^2 = Y$ and $x = X$, and obtain the complete primitive and singular solution, if any.

e) Solve the following differential equation by reducing to its normal form:

$$\frac{d^2y}{dx^2} - 4x \frac{dy}{dx} + (4x^2 - 3)y = e^{x^2}.$$

f) Solve the differential equation

$$\frac{d^3y}{dx^3} + y = e^{2x} \sin x + e^{\frac{x}{2}} \sin \frac{\sqrt{3}}{2} x.$$

11. Answer any **one** question: $10 \times 1 = 10$

a) i) Determine whether the equation

$$(1+yz)dx + x(z-x)dy - (1+xy)dz = 0$$

is integrable. Also obtain the integral of the equation, if integrable.

ii) If y_1 and y_2 are two linearly independent integrals of the differential equation

$$\frac{d^2y}{dx^2} + p_1 \frac{dy}{dx} + p_2 y = 0$$

where p_1, p_2 are functions of x , then show that

$$y_1 \frac{dy_2}{dx} - y_2 \frac{dy_1}{dx} = c e^{-\int p_1 dx}$$

where $c(\neq 0)$ is a constant. $4+6=10$

b) i) Reduce the differential equation

$$\sin^2 x \frac{dy}{dx} - 2y = 0 \text{ to exact form and hence}$$

solve it.

ii) Solve: $\frac{dx}{dt} + 5x + y = e^t$

$$\frac{dy}{dt} - x + 3y = e^{2t}. \quad 5+5=10$$
